

**TOPOLOGICAL PHASE TRANSITIONS
IN TWO-DIMENSIONAL SYSTEMS WITH
INTERNAL SYMMETRIES**

S.A.Bulgadaev

L.D.Landau Institute for Theoretical Physics
Kosyghin Str.2, Moscow, 117334, RUSSIA

A talk given at seminar

”Topological Defects in Non-Equilibrium Systems
and Condensed Matter”

Dresden, 22 July, 1999, Germany

CONTENT

I. Introduction.....	3
II. Multicomponent generalization and spaces with vector π_1	6
III. σ -models on T_G , duality and effective theories.....	10
IV. Phase diagram, critical properties and symmetries.....	13
V. d-dimensional conformal σ -model and topological excitations.....	19
VI. Conclusions.....	22
VII. References	22

I. Introduction

The topological phase transition (TPT) or the Berezinskii-Kosterlitz-Thouless (BKT) phase transition (PT) takes place in two-dimensional systems with degenerated vacuum, whose order parameter $\psi = e^{2\pi i\varphi} \in \mathcal{M} = S^1$. Among them are XY -model, superconductors, bose-liquids and many other systems.

The main unusual feathers of this TPT are:

- 1) absence of the spontaneous magnetization in low-T phase (Stanley, Kaplan 1966);
- 2) correlation functions in low-temperature phase must fall off algebraically (Rice 1966).

In particular, XY -model on a lattice is defined as follows

$$\mathcal{Z}_{XY} = \sum \exp(-\beta\mathcal{H}),$$

$$\mathcal{H} = -\frac{1}{2} \sum_{\langle i,j \rangle} J(\psi_i \psi_j^* + c.c.) = -J \sum_{\langle i,j \rangle} \cos(\varphi_i - \varphi_j).$$

Its continuous variant is a nonlinear σ -model (NSM) on S^1

$$\mathcal{Z}_{NS} = \int D\varphi e^{-\mathcal{S}[\varphi]}, \quad \mathcal{S}_{NS} = \frac{1}{2\alpha} \int |\partial\psi|^2 d^2x = \frac{1}{2\alpha} \int (\partial\varphi)^2 d^2x,$$

$$\alpha \simeq T/2J.$$

This form can be also considered as a longwave approximation of the Ginzburg-Landau (GL) theory

$$\mathcal{S}_{GL} = \int d^2x \left[\frac{1}{2\alpha} |\partial\psi|^2 + M^2(|\psi|^2 - 1)^2 \right]$$

with large M .

A circle S^1 has a discrete abelian first homotopic group π_1

$$\pi_1(S^1) = \mathbb{Z}.$$

Due to this property the topologically stable excitations, vortices, are possible in these systems.

One vortex solution has a form (at large distances)

$$\varphi(\mathbf{x}) = \frac{1}{\pi} \arctan \frac{y}{x} + \varphi_0, \quad \mathbf{x} = (x, y) \in \mathbb{R}^2,$$

where φ_0 is some constant. Its choice defines a form of vortex. For $\varphi_0 = 0$ it is pure potential, and for $\varphi_0 = \pi/2$ it is pure rotational. An account of vortices means that theory must be considered on the covering space $\mathbb{R}$ of the circle $S^1 = \mathbb{R}/\mathbb{Z}$.

The energy of one "vortex" with topological charge $e \in \mathbb{Z}$ is logarithmically divergent

$$E = \frac{e^2}{2\alpha} 2\pi \ln \frac{R}{a},$$

where R is a space radius, a is some short-wave cut-off parameter (a radius of the vortex core or a lattice constant).

The energy of N -"vortex" solution, E_N , with the full topological charge $e = \sum_{i=1}^N e_i = 0$ is finite and equals

$$E_N = \frac{2\pi}{2\alpha} \sum_{i \neq k}^N e_i e_k \ln \frac{|\mathbf{x}_i - \mathbf{x}_k|}{a} + C(a) \sum_i^N e_i^2,$$

here $C(a)$ is some nonuniversal constant, determining "self-energy" (or core energy) of vortices and depending on type of core regularization.

Berezinsky (1970,1971), Popov (1971,1976) and Kosterlitz and Thouless (1973,1974) have shown for the first time an important role of these topological excitations for existence of TPT in two-dimensional systems with continuous abelian symmetry.

These results have stimulated further investigations, given a discovery of monopoles (Polyakov 1974, t'Hooft 1974), instantons (Belavin et al. 1975), and others topological excitations in different branches of physics (Toulouse and Kleman 1976, Volovik and Mineev 1977).

The partition function $\mathcal{Z}_{XY}$ can be approximated by product of two partition functions (Kosterlitz 1974, Jose et al.1977)

$$\mathcal{Z}_{XY} \simeq \mathcal{Z}_{sw} \mathcal{Z}_{CG},$$

where $\mathcal{Z}_{CG}$ is the grand partition function of dilute Coulomb gas (CG) of topological excitations with minimal charges $e = \pm 1$ and $\mathcal{Z}_{sw}$ is the partition function of *free* "spin-waves". $\mathcal{Z}_{CG}$ can be represented, in its turn, in the

form of effective field theory with noncompact field ϕ and sine-Gordon action $\mathcal{S}_{SG}$ (Jose et al. 1977, Wiegmann 1978)

$$\mathcal{Z}_{CG} = \mathcal{Z}_{SG} = \int D\phi e^{-\mathcal{S}_{SG}[\phi]},$$

$$\mathcal{S}_{SG} = \int \left[\frac{1}{2\alpha} (\partial\phi)^2 - 2\mu^2 \cos \phi \right] d^2x$$

The TPT takes place in the system of vortices and can be described by the effective SG theory. This system has two different phases:

1) **high-T phase**: plasma-like, massive, with finite correlation length with essential singularity at T_c

$$\xi \sim a \exp(A\tau^{-\nu}), \quad \nu = \frac{1}{2},$$

$$\tau = \frac{T - T_c}{T_c} \rightarrow 0, \quad T_c \approx \pi J;$$

2) **low-T phase**: dielectric, massless, with infinite correlation length and algebraically falling correlations.

The TPT in low-dimensional systems can describe such important physical phenomena as compactification - decompactification (Green et al., 1988), localization - delocalization (Langer et al. 1987, Schmid 1983, Bulgadaev 1984) coherence - decoherence (Zaikin and Schon 1990) and others. For this reason it is very interesting to find the next

Possible generalizations of TPT on:

1) multicomponent 2D systems, having more complex vacuum manifolds $\mathcal{M}$ with nontrivial $\pi_1(\mathcal{M})$;

2) D-dimensional systems, admitting the interacting topological excitations with logarithmically divergent energies.

II. Multicomponent generalization and spaces with vector π_1

1. Usual tori.

The simplest generalization of circle

$$S^1 = e^{2\pi i \varphi}, \quad 0 \leq \varphi \leq 1$$

is a torus T^n , which equals to the direct product of n circles

$$T^n = \mathbb{R}^n / \mathbb{Z}^n = \bigotimes_{i=1}^n S_i^1,$$

where $\mathbb{R}^n$ is n -dimensional euclidean space, $\mathbb{Z}^n$ is n -dimensional simple cubic integer-valued lattice

$$\mathbb{Z}^n = \bigoplus_{i=1}^n \mathbb{Z}_i.$$

A torus T^n can appear as vacuum manifold of superposition of n independent abovementioned GL theories

$$\mathcal{S} = \sum_i^n \mathcal{S}_{GLi}$$


It is important to note, that a symmetry group of this theory is a semidirect product of T^n and a finite discrete group of permutations instead of usual continuous unitary or orthogonal groups.

The homotopy group of torus $\pi_1(T^n)$ is usually written as

$$\pi_1(T^n) = \bigoplus_{i=1}^n \mathbb{Z}_i = \mathbb{Z}^n,$$

where i -th component describes maps of the boundary S^1 into i -th circle of T^n . These circles form so called "bouquet" of circles (a set of circles with one common point)

$$B_n^1 = S_1^1 \vee \dots \vee S_n^1.$$



It is clear that maps into different components cannot be transformed into or annihilate each other. Consequently, one can introduce in $\pi_1(T^n)$ and in space of corresponding topological charges a vector structure. For this one needs:

- 1) a vector basis $\{\mathbf{e}_i\}$;
- 2) a metris or/and a scalar product in this space.

In this basis all topological charges must have integer-valued components, and a scalar product will determine interaction of different charges.

$$\mathbf{q} = \mathbf{q}_1 + \mathbf{q}_2, \quad q^2 = q_1^2 + q_2^2 + 2(\mathbf{q}_1 \mathbf{q}_2).$$

In general, this metrics is not fixed, but is defined by properties of the initial theory.

For T^n a basis $\{\mathbf{e}_i\}$ is the canonical one with

$$(\mathbf{e}_i \mathbf{e}_j) = \delta_{ik} = g_{ik}.$$

As a result the topological charges, corresponding to different S_i^1 , do not interact! Thus, the theories with $\mathcal{M} = T^n$ reduce from topological PT point of view to the case $\mathcal{M} = S^1$. Moreover, one can show that all tori $T_L = \mathbb{R}^n / \mathbb{L}$, where $\mathbb{L}$ is some full n -dimensional lattice in $\mathbb{R}^n$

$$\mathbb{L} = \sum_{i=1}^n n_i \mathbf{e}_i, \quad n_i \in \mathbb{Z}_i, \quad \mathbf{e}_i \in \{\mathbf{e}_i\}_L, \quad g_{ik} = \sum_{a=1}^n e_i^a e_k^a,$$

($\{\mathbf{e}_i\}_L, i = 1, \dots, n$ forms a basis of lattice $\mathbb{L}$) reduce in this sense to the case S^1 also. It means that for all systems, having such tori as a vacuum space, the TPT will have the same properties.

At the same time topological excitations on $\mathcal{M}$ with a rank of $\pi_1(\mathcal{M})$ $n \geq 2$ can have new additional properties. In particular, they can form knotted configurations in case $n = 2$. For example, a "trefoil" knot corresponds to $\mathbf{e} = (2, 3)$.

2. Degenerated (or constrained) tori.

For obtaining nontrivial generalizations one must to put on some constraints on tori T^n and/or on the initial theory. There is a regular method for obtaining such constraints (Bulgadaev 1996) which may be usefull in different regions of physics due to its relation only with theory of simple Lie group G .

Proposal:

One must consider theories connected with the Cartan tori T_G , the maximal abelian subgroups of the simple compact Lie groups G .

Tori T_G generalize a torus $T^n = T_{U(n)}$. They all can be considered as appropriately constrained torus T^N with large enough N . The corresponding GL type theories can be written out in abstract general form in terms of group G , but in the real physical systems their forms and corresponding symmetry breaking must be determined in each case separately, since the underlying group structure is usually hidden.

Two examples.

Let us represent a torus T^N in a form of diagonal matrix

$$T^n = \text{diag}(\exp(i\varphi_1), \dots, \exp(i\varphi_n)) = \\ \text{diag}(\exp(i(\mathbf{e}_1 \boldsymbol{\varphi})), \dots, \exp(i(\mathbf{e}_n \boldsymbol{\varphi}))),$$

where a vector $\boldsymbol{\varphi} = (\varphi_1, \dots, \varphi_n)$ is introduced.

Then

1) a condition $\sum_1^{n+1} \varphi_i = 0$ for torus T^{n+1} gives a torus of group $G = A_n = SU(n+1)$ in fundamental vector representation with vectors $\{\mathbf{e}_a\}$, $a = 1, \dots, n+1$, directed from origin to the vertices of hypertetrahedron in n -dimensional space;

2) a reduction of torus T^{2n} to the form $\varphi_i = -\varphi_{n+i}$, $i = 1, \dots, n$, giving an additional reflection symmetry $\varphi_i \rightarrow -\varphi_i$, $i = 1, \dots, n$, reproduces a torus of group $D_n = SO(2n)$ in fundamental vector representation with vectors $\{\mathbf{e}_a\}$, $a = 1, \dots, 2n$, directed to the vertices of n -dimensional hyperoctahedron.

In general, the Cartan torus T_G , the maximal abelian subgroup of group G , consists of elements

$$\mathbf{g} = e^{2\pi i(\mathbf{H} \boldsymbol{\phi})}, \quad \mathbf{H} = \{H_1, \dots, H_n\} \in \mathcal{C}, \quad [H_i, H_j] = 0,$$

where n is a rank of G , $\mathcal{C}$ is a maximal commutative Cartan subalgebra of the Lie algebra $\mathcal{G}$ of the group G . We assume here that $(\mathbf{H}\phi)$ is usual euclidean scalar product. Due to their commutativity, all H_i can be diagonalized simultaneously. Their eigenvalues, called the weights $\mathbf{w}$, depend on the concrete representation of G and $\mathcal{C}$. The weights $\{\mathbf{w}_a\}_\tau$, $a = 1, \dots, p$, belonging to a p -dimensional irreducible representation $\tau(G)$, form a set "of quantum numbers" of this representation.

All possible weights $\mathbf{w}$ of the simply connected group G (or of the universal covering group $\tilde{G}$ of non-simply connected group G) form a lattice of weights L_w . Its basis can be chosen in different ways. In some cases the more convenient basis is a basis of the vectors with minimal norms.

The weight lattice L_w , its reciprocal dual root lattice L_v , and related root lattice L_r are needed for finding out $\pi_1(T_G)$. The last can depend on the concrete representation $\tau(G)$. In two above examples the vectors directed to the vertices of regular polyhedra form the sets of weights of the corresponding fundamental representations, while their edges form the sets of roots (in this case the roots and dual roots coincide).

Since in any irreducible representation $\tau(G)$ of dimension p , one can choose the eigenvectors $|a\rangle$ of $\mathbf{H}$ as a basis

$$\mathbf{H}|a\rangle = \mathbf{w}_a|a\rangle, \quad a = 1, \dots, p,$$

then in this basis all H_i (and any element $\mathbf{g} \in T_G$) have diagonal form

$$\mathbf{g}_\tau = \text{diag}(e^{2\pi i(\mathbf{w}_1 \phi)}, \dots, e^{2\pi i(\mathbf{w}_p \phi)})$$

Here some weights can coincide, or even some weights can be equal to 0.

The main differences of this form from the usual representation of T_L type tori are:

1) a dimension of diagonal matrices coincides with dimension p of τ -representation, which is usually larger, than rank of G ;

2) the set of weights $\{\mathbf{w}\}_\tau$ has a discrete Weyl (or crystallographic) symmetry, which ensures Weyl invariance of T_G and results in the next two properties

$$\sum_{a=1}^p \mathbf{w}_a = 0, \quad g_{ik} = \sum_{a=1}^p w_i^a w_k^a = B_\tau \delta_{ik},$$

where constant B_τ depends on representation. In case of T_G the effective metrics g_{ik} is proportional to the euclidean one.

It is obvious, that in this representation all $\mathbf{g} \in T_G$ are periodic with a lattice of periods $\mathbb{L}_\tau^t$, inverse to the lattice L_τ , generated by weights $\mathbf{w}_a (a = 1, \dots, p)$ of τ -representation. It means that L_τ^t form a set of all topological charges of τ -representation of T_G .

The lattice L_τ^t satisfies the next restriction

$$L_{w^*} \supseteq L_\tau^t \supseteq L_v.$$

where L_{w^*} is a weight lattice of dual group G^* . For $\tau = \min$ a lattice $L_\tau^t = L_v$, for $\tau = \text{ad}$ a lattice $L_\tau^t = L_{w^*}$. The lattices L_v and L_{w^*} differ by a factor, which is isomorphic to the centre Z_G of group G

$$L_{w^*}/L_v = Z_G.$$

Thus the set $\{\mathbf{q}\}$ can vary from the set of minimal vectors (it forms the so called Voronoi region or Wigner - Seitz cell of the corresponding lattice) of the weight lattice till that of the root lattice. All possible cases are determined by subgroups of the centre Z_G . For groups G with $Z_G = 1$ the lattices L_v and L_{w^*} coincide.

When $L_\tau^t = L_w$ (or $L_\tau^t = L_{w^*}$) one can reproduce all weights (i.e. quantum numbers) of group G (or G^*) as the vector topological charges of vortices!

In this relation becomes obvious the important role of tori T_G , connected with self-dual lattices (they are $T_{U(n)}$ and T_{E_8}): all quantum numbers of corresponding groups have topological interpretation in terms of topological charges of vortices. When $L_\tau^t \in L_w$, one can enlarge L_τ^t by factorization of T_G over finite group $Z_\tau = L_w/L_\tau$.

III. σ -models on T_G , duality and effective theories

The euclidean two-dimensional nonlinear σ -models on T_G , generalizing σ -model on a circle S^1 , has the following form

$$\begin{aligned} \mathcal{S} &= \frac{1}{2\alpha} \int d^2x \text{Tr}_\tau (\mathbf{t}_\nu^{-1} \mathbf{t}_\nu) = \frac{(2\pi)^2}{2\alpha} \int d^2x \text{Tr}_\tau (\mathbf{H} \boldsymbol{\varphi}_\nu)^2 \\ &= \frac{(2\pi)^2}{2\alpha} B_\tau \int d^2x (\boldsymbol{\varphi}_\nu)^2, \end{aligned}$$

where $\varphi_\nu = \partial_\nu \varphi$, $\nu = 1, 2$. It will be convenient below to include a factor B_τ as a normalization factor into trace Tr_τ . This gives us a canonical euclidean metric in space of topological charges.

These theories are invariant under direct product of right (R) and left (L) groups $N_G^{R(L)}$, which are a semi-direct product of T_G and W_G

$$N_G = T_G \times W_G.$$

The group N_G is called a normalizator of T_G and is a symmetry group of torus T_G .

NSM on T_G are the multicomponent generalization of XY -model, having properties analogous to those of XY -model:

- 1) a zero beta-functions $\beta(\alpha)$ due to flatness of T_G ;
- 2) non-trivial homotopy group π_1 and corresponding vortex solutions.

The general N -vortex solutions have the next form

$$\varphi(\mathbf{x}) = \sum_{i=1}^N \mathbf{q}_i \frac{1}{\pi} \arctan\left(\frac{y - y_i}{x - x_i}\right),$$

$$\mathbf{q}_i \in L_\tau^t, \quad (\mathbf{q}_i \mathbf{w}_a) \in \mathbb{Z}, \quad (x, y) \in \mathbb{R}^2.$$

Here L_τ^t is a lattice of all possible topological charges in τ -representation. The energy of N -vortex solution, E_N , with a whole topological charge $\sum_{i=1}^N \mathbf{q}_i = 0$ is

$$E_N = \sum_i E_{q_i}^0 + E_{Nint}, \quad E_{q_i}^0 = \frac{1}{2\alpha} C(a)(\mathbf{q}_i \mathbf{q}_i),$$

$$E_{Nint} = \frac{2\pi}{2\alpha} \sum_{i \neq k}^N (\mathbf{q}_i \mathbf{q}_k) \ln \frac{|x_i - x_k|}{a},$$

where $E_{q_i}^0$ is a "self-energy" (or an energy of the core) of vortex with topological charge $\mathbf{q}_i$. Only such solutions give finite contribution to partition function of the theory.

Analogous topological excitations with vectorial charges exist also in the models of two-dimensional crystall melting (Nelson 1978).

In quasi-classical approximation (or in low T expansion) one can represent a partition function of the σ -model on T_G

$$\mathcal{Z} = \int D\varphi \exp(-\mathcal{S}[\varphi])$$

as a grand partition function of classical neutral Coulomb gas of vortex solutions with minimal isovectorial topological charges $\mathbf{q}_i \in \{\mathbf{q}\}_\tau$, where $\{\mathbf{q}\}_\tau$ is a set of minimal vectors of lattice of all possible topological charges in τ -representation L_τ^t

$$\mathcal{Z} = \mathcal{Z}_0 \mathcal{Z}_{CG}, \quad \mathcal{Z}_{CG} = \sum_{N=0}^{\infty} \frac{\mu^{2N}}{N!} \sum'_{\{\mathbf{q}\}} \mathcal{Z}_N(\{\mathbf{q}\}|\beta).$$

Here $\mathcal{Z}_0$ is a partition function of free massless isovectorial boson field which corresponds to "spin waves" of XY -model

$$\mathcal{Z}_0 = \int D\phi \exp(-\mathcal{S}_0[\phi]),$$

$\sum'$ in $\mathcal{Z}_{CG}$ goes over all neutral configurations of minimal charges $\mathbf{q}_i \in \{\mathbf{q}\}_\tau$ with $\sum_1^N \mathbf{q}_i = 0$,

$$\mathcal{Z}_N(\{\mathbf{q}\}|\beta) = \prod_{i=1}^N \int d^2x_i \exp(-\beta H_N(\{\mathbf{q}\}))$$

$$H_N(\{\mathbf{q}\}) = \sum_{i<j}^N (\mathbf{q}_i \mathbf{q}_j) D(x_i - x_j),$$

$$D(x) = \int \frac{d^2k}{(2\pi)^2} (e^{i(\mathbf{k}\mathbf{x})} - 1) f(ka)/k^2 \underset{|x| \gg a}{\sim} \frac{1}{2\pi} \ln |x/a|$$

where

$$\mu^2 = a^{-2} y^2 (det)^{-1/2}, \quad y^2 = e^{-E_q^0}$$

is a chemical activity of Coulomb gas, det is a determinant of the fluctuations over one vortex solution (further we will suppose that it is equal to some constant of order $O(1)$ and assume that $det = 1$),

$$\beta = 4(\pi)^2/\alpha,$$

$f(ka)$ is a regularisator such that

$$\lim_{k \rightarrow 0} f(ka) = 1, \quad \lim_{k \rightarrow \infty} f(ka) = 0.$$

There is a connection between compact σ -models on T_G and noncompact generalized SG theories

$$\mathcal{Z}_{CG} = \int D\phi e^{-\mathcal{S}_{eff}}, \quad \mathcal{S}_{eff} = \int \frac{1}{2\beta} (\partial\phi)^2 - \mu^2 V(\phi),$$

$$V(\phi) = \sum_{\{\mathbf{q}\}} \exp i(\mathbf{q}\phi).$$

where $\sum_{\{\mathbf{q}\}}$ goes over the set of minimal topological charges, and $\phi \in \mathbb{R}^n$. Strictly speaking, the effective theories with arbitrary parameters μ and β are more general than initial σ -models.

The account of vortices reduces the initial symmetry group N_G into discrete dual group $W_{G^*} \times L_q^{-1}$ (W_{G^*} is a dual Weyl group, L_q^{-1} is a periodicity lattice of potential V). This dual group generalizes the dual group $Z_2 \times \mathbb{Z}$ of XY -model.

Thus, in this semiclassical and long wavelength approximation

Compact theory on a torus T_G with *continuous* symmetry N_G appears equivalent (again modulo $\mathcal{Z}_0$) to *noncompact* theory with periodic potential and an *infinite discrete* symmetry.

For simply laced groups the potentials V for groups from series A, D, E ($A_n = SU(n+1), D_n = SO(2n), E = E_{6,7,8}$) coincide with characters of adjoint representations of these groups (modulo some constant, corresponding to zero weight). In this case the general theories can describe systems with symmetry G broken to N_G .

The noncompact theories can be considered also as corresponding linear σ -models. For this reason

Compact nonlinear σ -models on T_G are equivalent in this approximation to *noncompact linear* σ -models on Cartan tori of dual group T_{G^*} .

IV. Phase diagram, critical properties and symmetries

The BKT type PT can be investigated by renormalization of the corresponding effective field theories (Wiegmann 1978, Bulgadaev 1981,1996a).

Under renormalization transformations both parameters μ and β are renormalized. It is convenient to introduce two dimensionless parameters

$$(\mu a)^2 = g, \quad \delta = \frac{\beta q^2 - 8\pi}{8\pi}$$

where q^2 is a square of the norm of the minimal vector topological charges from $\{\mathbf{q}\}$. The effective theories are renormalizable only if the vectors from $\{\mathbf{q}\}$ belong to some lattice (here L_τ^t). A new critical properties can appear only if a geometry of $\{\mathbf{q}\}$ is such that each vector $\mathbf{q} \in \{\mathbf{q}\}$ can be represented as a sum of two other vectors from $\{\mathbf{q}\}$. It means that a potential V has the next property

$$V^2 = \sum_a \sum_b \exp(i(\mathbf{q}_a + \mathbf{q}_b, \phi)) = \theta_0 + \theta V + \sum_{a+b \neq c} \exp(i(\mathbf{q}_a + \mathbf{q}_b, \phi)),$$

where θ_0 is doubled number of vectors in $\{\mathbf{q}\}$, having their opposite vectors, θ is a number of times how each vector $\mathbf{q}$ can be represented as a sum of two other vectors from $\{\mathbf{q}\}$, the last term contains vectors with norm larger than $|q|$ (so called higher harmonics). An appearance of V on the right hand is very restrictive (the angles between vectors can be equal only to $\pi/3, 2\pi/3$ or $\pi/2$) and coincides with a definition of the root systems $\{\mathbf{r}\}$ of simple groups from series A, D, E or with a definition of the root set of the even integer-valued (in some scale) lattices of $\mathbb{A}, \mathbb{D}, \mathbb{E}$ types. The sets of minimal roots (or minimal dual roots) of all simple groups belong to four series of integer-valued lattices $\mathbb{A}, \mathbb{D}, \mathbb{E}, \mathbb{Z}$. For theories with sets $\{\mathbf{q}\} \notin \mathbb{A}, \mathbb{D}, \mathbb{E}$, all critical properties will be the same as for theories with $\{\mathbf{q}\} \in \mathbb{Z}^n$.

The RG equations for $\{\mathbf{q}\}$ from these (i.e. $G = A, D, E$) lattices have the next form

$$\frac{dg}{dl} = -2\delta g + B_G g^2, \quad \frac{d\delta}{dl} = -C_G g^2.$$

Here $B_G = \pi\theta_G$, and $C_G = 2\pi K_G$, where K_G is the value of the second Casimir operator in adjoint representation (where $\mathbf{w}_a = \mathbf{r}_a$, the roots)

$$\sum_a r_i^a r_j^a = K_G \delta_{ij}.$$

The RG equations of such type with coefficients corresponding to the case $G = A_2$ have been obtained firstly by Nelson (1978) under investigation of the melting of the two-dimensional triangle lattice.

The eigenvalue of the second Casimir operator K_G for groups from series A, D, E can be expressed in terms of the corresponding Coxeter number h_G

$$K_G = 2h_G, \quad h_G = \frac{(\text{number of all roots})}{(\text{rank of group})},$$

$$h_{A_n} = n + 1, \quad h_{D_n} = 2(n - 1), \quad h_{E_{6,7,8}} = 12, 18, 30.$$


This definition of the Coxeter number coincides with that of the Coxeter number of the corresponding integer-valued lattices from series $\mathbb{A}, \mathbb{D}, \mathbb{E}$. The coefficient B_G can be expressed also through the Coxeter number

$$\theta_G = K_G - 4 = 2(h_G - 2).$$

Thus, all coefficients of RG equations depend only on the Coxeter number h_G or on the second Casimir value K_G .

The RG equations have two separatrices with next declinations (u_1 corresponds to the phase separation line).

$$u_{1,2} = (1/\pi h_G, -1/2\pi).$$



Schematic phase diagram.

The dashed line of the initial values corresponds to the initial σ -model. This line is defined by the dependencies of the parameters β and μ on coupling constant α .

The critical exponent ν_G , determining divergence of the correlation length ξ

$$\xi \sim a \exp(A\tau^{-\nu_G}), \quad \tau = \frac{T - T_c}{T_c},$$

is the inverse of the Lyapunov index of the separatrix 1 and equals

$$\nu_G = 2/(2 + h_G).$$

Since h_G is a relatively rough characteristic, ν_G and other critical properties of different systems can coincide.

All possible values of exponent ν_G are the next (Bulgadaev 1983,1996a)

G	A_n	B_n	C_n	D_n	G_2	F_4	E_6	E_7	E_8
ν_G	$\frac{2}{n+3}$	$\frac{1}{n}$	$\frac{1}{2}$	$\frac{1}{n}$	$\frac{2}{5}$	$\frac{1}{4}$	$\frac{1}{7}$	$\frac{1}{10}$	$\frac{1}{16}$

The series A_n gives a maximal set of possible values of ν_G : $1/k$ and $2/(2k+1)$ (where k are integers ≥ 2).

Low-T phase properties

In the low-temperature phase the correlation functions of the fields exponentials equal to the free correlation functions with a renormalized "temperature" $\bar{\beta}$ which depends on initial values β_0 , $\bar{\beta} = \lim_{l \rightarrow \infty} \beta(l)$:

$$\left\langle \prod_{s=1}^t \exp(i(\mathbf{r}_s \phi(x_s))) \right\rangle = \prod_{i \neq j}^t \left| \frac{x_i - x_j}{a} \right|^{\bar{\beta}(\mathbf{r}_i \mathbf{r}_j)/2\pi},$$

$$\sum_{i=1}^t \mathbf{r} = 0.$$

At the PT point (where $\bar{\beta} = \beta^* = 8\pi/r^2 = 4\pi$) an additional logarithmic factor, related with the "null charge" behaviour of g and δ on the critical separatrix (the phase separation line), appears in them:

$$\prod_{i \neq j}^t \left(\ln \left| \frac{x_i - x_j}{a} \right| \right)^{\beta^*(\mathbf{r}_i \mathbf{r}_j)/2\pi A_G} = \prod_{i \neq j}^t \left(\ln \left| \frac{x_i - x_j}{a} \right| \right)^{h_G \cos(\mathbf{r}_i \mathbf{r}_j)},$$

where $A_G = 4/h_G$ is a coefficient in RG equations for δ on the critical separatrix 1.

In this phase additional vortices can move like particles in dielectric media.

Free-like behaviour of the low-temperature phase (except logarithmic corrections at criticality) gives a possibility to use for their description conformal field theories with integer central charges $C = n$, instead of PT points of two-dimensional systems with discrete symmetries, which are described by conformal theories with rational central charges (Belavin et al. 1984, Dotsenko and Fateev 1984,1985). The BKT PT can be considered as some degeneration of II order PT: it corresponds to the limiting case $k \rightarrow \infty$ (where k is a level) of the sequence of the restricted SG models, describing minimal conformal theories (Smirnov and Reshetikhin 1989, LeClair 1989). Analogously, TPT in σ -models on T_G are the limiting cases of unitary minimal conformal theories, connected with conformal W -algebras (Fateev and Lukyanov 1988). In this relation it is interesting to note that there exists a puzzling coincidence of ν_G with "screening" factor in formulas for central charges of the affine Lie algebras $\hat{\mathfrak{g}}$ (Kac 1990) at level $k = 2$ (though T_G corresponds to $k = 1$)

$$C_k = \frac{k}{k + h_G} \dim G$$

and of coset realization of the minimal unitary conformal models (Goddard et al. 1985,1986) at level $k = 1$

$$C_k = r \left(1 - \frac{h_G(h_G + 1)}{(k + h_G)(k + h_G + 1)} \right).$$

Properties of massive phase

In this phase all additional vector charges will be shielded like in plasma.

A separatrix 2 is a boundary of the asymptotically free (AF) region in UV-limit. There is also another possibility of the enhancement of the symmetry of the initial nonlinear σ -model on this separatrix. σ -model has at classical level two symmetries: 1) scale (or conformal) symmetry, 2) isotopic global symmetry $N_G = T_G \times W_G$. Conformal symmetry is spontaneously broken in IR region by vortices. For this reason σ -model has in massive phase a finite correlation length $\xi \sim m^{-1}$, where m is a characteristic mass scale of the theory. This mass must depend on the coupling constant α or β . The behaviour of m near PT point is described by above formula, where

$$\tau \sim \frac{\alpha - \alpha_c}{\alpha_c}.$$

On separatrix 2 one obtains

$$m \sim \Lambda \exp(-1/2\pi g h_G), \quad \Lambda \sim a^{-1}.$$

The fact that on a separatrix 2 a mass scale is defined by $K_G \sim h_G$ is a corollary of the independence of its declination on G .

This expression coincides with those for chiral models on groups (Polyakov 1975, Ogievetskii, Wiegmann, 1986) and with those, obtained from an exact solution of the appropriate fermion theories in main approximation on g (Destri, De Vega 1987)

$$m \sim \Lambda \exp(-2\pi/(gK_G/2)).$$

Thus,

Mass scale on separatrix 2 coincides (at least for $G = A, D, E$) with that for all G -invariant theories (chiral and fermionic), connected with simple Lie groups G .

It means a possible restoration of the full symmetry group G in σ -models with reduced symmetries N_G . This can be seen also from the equivalence of the effective field theories in cases $G = A_{n-1} = SU(n), G = D_n = O(2n)$, to the fermion theories with the same global symmetry groups G (Coleman 1975, Mandelstam 1975, Bulgadaev 1981, 1983)

There is strong dependence of the index $\nu(\alpha)$, which interpolates between $\nu = \nu_G$ near the PT point and $\nu = 1$ near the separatrix 2. The first region corresponds to symmetry of T_G , which is a torus normalisator $N_G = T_G \times W_G$, while the second one corresponds to more symmetric, G -invariant, situation.

Another interesting application of TPT in NSM on T_G is connected with string theory and its cosmological aspects. In this case TPT can be considered as effective decompactification PT (Bulgadaev 1998), a massive phase corresponds to a compactified phase and a massless phase corresponds to a decompactified phase.

V. d-dimensional conformal σ -model and topological excitations

1. d-dimensional generalization.

Importance of topological excitations with logarithmically divergent energies in 1d systems has been earlier discovered by Anderson, Yuval and Hamann in 1970, in papers devoted to the Kondo problem. It was shown there that PT takes place in 1d Ising system with long-range interaction

$$J(r) \sim 1/r^2$$

due to the presence of logarithmically interacting domain walls.

The next properties unify these two models and give the conditions of existence of topological excitations with logarithmic energy in d-dimensional NS models:

- 1) conformal invariance at classical level,
- 2) homotopical group $\pi_{d-1}(\mathcal{M})$ must be nontrivial abelian discrete.

The first property defines form of action S and the second one defines partially a topology of $\mathcal{M}$

$$S = \frac{1}{2\alpha} \int d^d x d^d x' \psi_a(x) \boxtimes_{ab}^{(d)}(x - x') \psi_b(x'),$$

where $\psi \in \mathcal{M}$, $a, b = 1, 2, \dots, n$, n is a dimension of $\mathcal{M}$ and a form of kernel $\boxtimes$ depends on dimension of space d . For decoupled internal and physical spaces $\boxtimes$ can be decomposed

$$\boxtimes_{ab}^{(d)}(x) = g_{ab} \square_d(x),$$

where g_{ab} is the Euclidean metric of the space $\mathbb{R}^{N(n)}$, in which a manifold $\mathcal{M}$ can be embedded. In the momentum space, for small k

$$\square_d(k) \simeq |k|^d (1 + a_1(ka) + \dots),$$

where a is a UV cut-off parameter. Action $\mathcal{S}$ can be named d -dimensional *conformal* nonlinear σ -model. The kernel $\square_d$ generalizes an usual local and conformal kernel of two-dimensional σ -model

$$\square(k) \equiv \square_2(k) = k^2.$$

For local models an expression for $\boxtimes$ can be defined in terms of manifold $\mathcal{M}$ only

$$\boxtimes(x) = g_{ab}(\phi) \square \delta(x)$$

In odd dimensions $\square_d$ is nonlocal

$$\square_d(x) \sim 1/|x|^{2d},$$

but the models with such kernels are used often in physics.

The simplest spaces with such properties are the spheres S^{d-1} . In 3d case this is a conformal (or van der Waals, since $\square_3(x) \sim 1/|x|^6$) NS-models on sphere S^2 .

An existence of topological excitations with logarithmic energy follows from the invariance of the kernel B and the simplest topologically nontrivial excitations $\mathbf{n}_i = x_i/r$ under scale and conformal transformations

$$x_i \rightarrow x'_i = x_i/r^2, \quad r \rightarrow r' = 1/r, \quad x_i/r = x'_i/r',$$

$$d^d x \rightarrow d^d x / |\mathbf{x}|^d, \quad \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|^{2d}} \rightarrow \frac{|\mathbf{x}_1|^d |\mathbf{x}_2|^d}{|\mathbf{x}_1 - \mathbf{x}_2|^{2d}},$$

and, consequently,

$$\mathcal{S} \sim \int d^d x_1 d^d x_2 \frac{(\mathbf{x}_1 \mathbf{x}_2)}{r_1 r_2} \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|^{2d}}$$

is invariant and dimensionless. By direct calculation one can show that

$$\mathcal{S}[\mathbf{n}] \approx C_d \ln \frac{R}{a}$$

where $C_d = \frac{16}{\alpha}$, and different excitations interact through potential $G(r)$, inverse to $\square_3(x)$, which has a logarithmic behaviour at large distances

$$G(r) \sim \ln \frac{r}{R}.$$

2. Multicomponent generalization for $d > 2$.

A simple generalization of the sphere S^{d-1} , analogous to the torus T^n in 2d case, is a bouquet of n spheres

$$B_n^{d-1} = S_1^{d-1} \vee \dots \vee S_n^{d-1}$$

and all spaces $\mathcal{M}$ with this first nontrivial topological cell complex. But the corresponding vector topological charges will not interact as in a case of torus. For obtaining an interaction of topological charges in 3d case one needs to consider NS-models on deformed B^n . In particular, NS-models on the maximal flag spaces $F_G = G/T_G$ of the simple compact groups G , with $\pi_2(F_G) = \mathbb{L}_v$, (note that a sphere S^2 is a particular case of F_G : $S^2 = SU(2)/U(1)$) will also have topological excitations with interacting vector topological charges $\mathbf{Q} \in \mathbb{L}_v$ and logarithmic energy.

Thus, these excitations will have the mixed properties of the two-dimensional vortices and instantons: their topology is connected with π_2 , but they interact as vortices.

VI. Conclusions

1. It is shown that one must consider deformed tori for obtaining the interacting vector topological charges.
2. Vector topological charges form a lattice and in some cases can reproduce all quantum numbers of the corresponding groups.
3. A sequence of approximately equivalent transformations of 2d models is constructed

General GL Theory $\rightarrow$ NS Model $\rightarrow$ TopExcGas $\rightarrow$ General SG Theory

It simplifies a problem and extracts all necessary long-wave properties of these theories! Here last theory has a pure group-theoretical structure and is universal for whole class of theories with the same symmetries.

4. All critical properties are classified by integer-valued lattices from series $\mathbb{A}, \mathbb{D}, \mathbb{E}, \mathbb{Z}$ and are characterised by the corresponding Coxeter numbers.
5. The possible scenarios of the dynamical enlargening of the initial internal symmetry groups is proposed.
6. D-dimensional conformal NS-models, having the topological excitations with logarithmic energies, are constructed.

I would like to thank the organizers of the workshop for the invitation and the opportunity to give this talk.

References

- [1] Anderson P.W., Yuval G., Hamann D.R., Phys.Rev. **B1** (1970) 4464.
- [2] Belavin A.A., Polyakov A.M., Schwartz A.S., Tyupkin Yu.S. **59B** (1975) 85.
- [3] Belavin A.A., Polyakov A.M., Zamolodchikov A.B., Nucl.Phys. **B241** (1984) 333.
- [4] Berezinskii V.L., JETP **59** (1970) 907, **61** (1971) 1144.
- [5] Bulgadaev S.A., Phys.Lett. **86A** (1981) 213; Theor.Math.Phys. **49** (1981) 7.
- [6] Bulgadaev S.A., Nucl.Phys. **B224** (1983) 349.
- [7] Bulgadaev S.A., JETP Letters **39** (1984) 264.
- [8] Bulgadaev S.A., JETP Letters **63** (1996)a 780; hep-th/9906091.
- [9] Bulgadaev S.A., JETP Letters **63** (1996) 796; hep-th/9901035.
- [10] Bulgadaev S.A., hep-th/9811226, (1998).
- [11] Coleman S., Phys.Rev., **D11** (1975) 2088.
- [12] Destri C., de Vega H.J., Preprint CERN-TH, 4895/87.
- [13] Dotsenko V.I., Fateev V.A., Nucl.Phys. **B240** (1984) 312, **B251** (1985) 691.

- [14] Fateev V.A., Lukyanov S.L., Int.Jour.Mod.Phys. **A13** (1988) 507.
- [15] Goddard P., Kent A., Olive D., Phys.Lett.**B152** (1985) 88, Commun.Math.Phys. **103** (1986) 105.
- [16] Green M., Schwarz J.H., Witten E., Superstrings Theory, Cambridge, 1988, vol.1,2.
- [17] t'Hooft G., Nucl.Phys.**B79** (1974) 276.
- [18] Jose J., Kadanoff L., Kirkpatrick S., Nelson D., Phys.Rev. **B16** (1977) 1217.
- [19] Kac V.N., Infinite dimensional Lie algebras, Cambridge University Press, 1990.
- [20] Kosterlitz J.M., Thouless J.P., J.Phys.**C6** (1973) 118.
- [21] Kosterlitz J.M., J.Phys.**C7** (1974) 1046.
- [22] LeClair A., Preprint PUPT-1124 (1989).
- [23] Legget A.J. et al. Rev.Mod.Phys. **59** (1987) 1.
- [24] Mandelstam S., Phys.Rev. **D11** (1975) 3026.
- [25] Nelson D.R., Phys.Rev. **B18** (1978) 2318.
- [26] Ogievetskii E., Wiegmann P.B., Phys.Lett. **168B** (1986) 360.
- [27] Polyakov A.M., Pisma v ZETP **20** (1974) 430.
- [28] Polyakov A.M., Gauge Fields and Strings, Harwood Academic Publishers, 1987.
- [29] Popov V.N., unpublished (1971); Feynman integrals in quantum field theory and statistical mechanics, Moscow. Atomizdat. 1976.
- [30] Reshetikhin N., Smirnov F., Commun.Math.Phys., **31** (1990) 157.
- [31] Rice T.M., Phys.Rev.**140** (1966) 1889.
- [32] Schmid A., Phys.Rev.Lett. **51** (1983) 1506.

- [33] Schon G., Zaikin A.D., Phys.Rep. **198** (1990) 237.
- [34] Stanley H.E., Kaplan T.A., Phys.Rev.Lett., **17** (1966) 913.
- [35] Toulouse G., Kleman M., J.Physique Lett. **37** (1976) 149.
- [36] Volovik G.E., Mineev V.P., JETP **45** (1977) 1186
- [37] Wiegmann P.B., J.Phys.**C11** (1978) 1583.